
How to Specify a Reference Point in Hypervolume Calculation for Fair Performance Comparison

Hisao Ishibuchi

hisao@sustc.edu.cn

Shenzhen Key Laboratory of Computational Intelligence, Department of Computer Science and Engineering, Southern University of Science and Technology, Shenzhen, 518005, China

Ryo Imada

ryo.imada@ci.cs.osakafu-u.ac.jp

Department of Computer Science and Intelligent Systems, Osaka Prefecture University, Sakai, 5998531, Japan

Yu Setoguchi

yu.setoguchi@ci.cs.osakafu-u.ac.jp

Department of Computer Science and Intelligent Systems, Osaka Prefecture University, Sakai, 5998531, Japan

Yusuke Nojima

nojima@cs.osakafu-u.ac.jp

Department of Computer Science and Intelligent Systems, Osaka Prefecture University, Sakai, 5998531, Japan

Abstract

The hypervolume indicator has frequently been used for comparing evolutionary multi-objective optimization (EMO) algorithms. A reference point is needed for hypervolume calculation. However, its specification has not been discussed in detail from a viewpoint of fair performance comparison. A slightly worse point than the nadir point is usually used for hypervolume calculation in the EMO community. In this paper, we propose a reference point specification method for fair performance comparison of EMO algorithms. First, we discuss the relation between the reference point specification and the optimal distribution of solutions for hypervolume maximization. It is demonstrated that the optimal distribution of solutions strongly depends on the location of the reference point when a multi-objective problem has an inverted triangular Pareto front. Next, we propose a reference point specification method based on theoretical discussions on the optimal distribution of solutions. The basic idea is to specify the reference point so that a set of well-distributed solutions over the entire linear Pareto front has a large hypervolume and all solutions in such a solution set have similar hypervolume contributions. Then, we examine whether the proposed method can appropriately specify the reference point through computational experiments on various test problems. Finally, we examine the usefulness of the proposed method in a hypervolume-based EMO algorithm. Our discussions and experimental results clearly show that a slightly worse point than the nadir point is not always appropriate for performance comparison of EMO algorithms.

Keywords

Evolutionary multi-objective optimization, hypervolume, reference point, performance comparison.

1 Introduction

In the field of evolutionary multi-objective optimization (EMO), a number of new algorithms continue to be proposed every year. The proposed algorithms are compared with existing ones using performance indicators. The hypervolume indicator (Zitzler and Thiele (1998)), which is also abbreviated as “hypervolume” in this paper, has frequently been used for performance comparison in the EMO community while a wide variety of indicators are available in the literature (Zitzler et al. (2003)). This is because no other Pareto compliant indicator is known (Zitzler et al. (2007)). The hypervolume indicator has also frequently been used in indicator-based EMO algorithms such as LAHC (Knowles et al. (2003)), SMS-EMOA (Emmerich et al. (2005); Beume et al. (2007)) and HypE (Bader and Zitzler (2011)). In these algorithms, multi-objective optimization is handled as single-objective optimization of a solution set for hypervolume maximization.

An important research issue is the analysis of the optimal distribution of solutions for hypervolume maximization. Theoretical studies (Emmerich et al. (2007); Auger et al. (2012); Brockhoff (2010); Friedrich et al. (2015)) show that the hypervolume indicator is maximized by a set of uniformly distributed solutions on a linear Pareto front of a two-objective problem. It is also shown that such a solution set is not optimal when the Pareto front is nonlinear. This is because the optimal distribution of solutions depends on the slope of the Pareto front in the two-dimensional objective space. Except for the case of single-dimensional degenerate Pareto fronts (Shukla et al. (2014)), no optimal distribution of solutions has been theoretically derived for multi-objective problems with three or more objectives. General theoretical analysis of the optimal distribution of solutions for hypervolume maximization is difficult for multi-objective problems with three or more objectives.

Selection of a pre-specified number of solutions from a given solution set has been actively studied for hypervolume maximization under the name of hypervolume subset selection. This is a single-objective combinatorial optimization problem. Efficient subset selection methods have been proposed in the literature (Basseur et al. (2016); Bringmann et al. (2014); Guerreiro et al. (2015, 2016); Kuhn et al. (2016)). Hypervolume subset selection can be also viewed as a pre-processing procedure for selecting a small number of solutions to be presented to the decision maker from a large number of solutions obtained by EMO algorithms (Ishibuchi et al. (2014)). Recently, such a pre-processing procedure has been used for performance comparison of EMO algorithms with an unbounded archive population (Ishibuchi et al. (2016); Tanabe et al. (2017)).

A reference point is needed to calculate the hypervolume indicator of a given solution set. For two-objective problems, the effect of the reference point specification on the optimal distribution of solutions for hypervolume maximization has been theoretically studied (Auger et al. (2012); Brockhoff (2010); Friedrich et al. (2015)). For three-objective problems with linear Pareto fronts, the optimal distribution of solutions has been empirically studied through a number of computation experiments (Ishibuchi et al. (2017a)). It is also shown through computational experiments that performance comparison results among different EMO algorithms depend on the location of the reference point (Ishibuchi et al. (2015, 2017c)). However, to the best of our knowledge, the reference point specification has not been studied in detail from a viewpoint of

This article is an extended version of two conference papers (Ishibuchi et al. (2017b)), which includes many new experiments, explanations and discussions on the reference point specification for multi-objective problems with nonlinear Pareto fronts and also for many-objective problems. The visualization method of hypervolume contributions is also improved from the conference paper (Ishibuchi et al. (2017b)).

fair performance comparison of EMO algorithms, especially for many-objective problems, except for our former conference paper (Ishibuchi et al. (2017b)) where it was discussed for two multi-objective test problems with linear Pareto fronts (i.e., DTLZ1 and inverted DTLZ1). This paper is an extended version of our former conference paper. We discuss the reference point specification not only for multi-objective test problems with linear Pareto fronts but also for those with non-linear Pareto fronts (i.e., DTLZ2 and Minus-DTLZ2), a multi-objective knapsack problem and a multi-objective car-side impact problem. A visualization method of hypervolume contributions is improved from the conference paper. Moreover, detailed discussions on the reference point specification in SMS-EMOA (Emmerich et al. (2005); Beume et al. (2007)) are added.

A general guideline for the reference point specification is to use a slightly worse point than the nadir point so that the reference point is dominated by all Pareto optimal solutions (i.e., so that all Pareto optimal solutions in a solution set have positive hypervolume contributions). However, there exists no specific numerical guideline. As a result, different specifications were used in different studies as pointed out by Chugh et al. (2018). For example, the reference point was specified for DTLZ1 (Deb et al. (2002)) with the ideal point $(0.0, \dots, 0.0)$ and the nadir point $(0.5, \dots, 0.5)$ as follows: 1% larger than the nadir point, i.e., $(0.505, \dots, 0.505)$ in Seada and Deb (2016), 10% larger in Yuan et al. (2016a,b) and Maltese et al. (2018), 40% larger in Wagner et al. (2007), and 100% larger in Li et al. (2015). In Zhang et al. (2018), the nadir point was used as the reference point. These observations raise the following two questions: “How sensitive are the performance comparison results to the reference point specification?” and “How should we specify the reference point for fair performance comparison?”.

The goal of this paper is to answer these two questions. First, we discuss the optimal distribution of solutions for hypervolume maximization. We clearly demonstrate that the sensitivity of the optimal distribution of solutions to the location of the reference point is problem-dependent. The effect of the reference point is minor when a multi-objective problem has a triangular Pareto front such as DTLZ1-4 (Deb et al. (2002)) and WFG4-9 (Huband et al. (2006)). However, the reference point specification has a large effect on the optimal distribution of solutions when a multi-objective problem has an inverted triangular Pareto front such as Minus-DTLZ1-4 and Minus-WFG4-9 (Ishibuchi et al. (2017c)). We discuss why the reference point specification has a different effect depending on the shape of the Pareto front. Then, we propose a reference point specification method for fair performance comparison of EMO algorithms for multi-objective and many-objective problems. Our idea is based on the following two assumptions with respect to the performance evaluation of solution sets by the hypervolume indicator.

1. A uniformly distributed solution set over an entire linear Pareto front should be highly evaluated by the hypervolume indicator. That is, such a solution set should have a large hypervolume.
2. All solutions from such a solution set should have similar hypervolume contributions. That is, no solution should have a dominant or too small effect on hypervolume calculation.

This paper is organized as follows. In Section 2, we examine the optimal distribution of solutions for hypervolume maximization using SMS-EMOA (Emmerich et al. (2005); Beume et al. (2007)) for DTLZ1-4 and Minus-DTLZ1-4. It is clearly demonstrated that the optimal distribution of solutions strongly depends on the location of the reference point when a multi-objective problem has an inverted triangular Pareto front. In

Section 3, we propose a reference point specification method for fair performance comparison based on the above-mentioned two assumptions and the theoretical studies on the optimal distribution of solutions for a two-objective problem with a linear Pareto front. In Section 4, we examine whether the proposed method can appropriately specify a reference point for fair performance comparison of EMO algorithms through computational experiments on various test problems. In Section 5, we examine the usefulness of the proposed method in SMS-EMOA through computational experiments. Finally, we conclude this paper in Section 6.

2 Optimal Distribution of Solutions for Hypervolume Maximization: Empirical Discussions

In the EMO community, we implicitly assume that a set of well-distributed solutions over the entire Pareto front of a multi-objective problem is a good solution set. We also implicitly assume that such a solution set is highly evaluated by the hypervolume indicator and the IGD (inverted generational distance indicator). Thus these two indicators have frequently been used for performance comparison in the EMO community. However, we do not know the optimal distribution of solutions with respect to each indicator except for some special cases such as two-objective problems. That is, we are comparing different EMO algorithms using these indicators without knowing which solution set of a fixed size (what type of solution sets of a fixed size) is most highly evaluated by each indicator. If a set of solutions on a particular region of the Pareto front is evaluated as being better than a set of well-distributed solutions over the entire Pareto front, performance comparison results may be inconsistent with our intuition (i.e., they may be unreliable). In this sense, it is important to know the optimal distribution of solutions with respect to each indicator. Actually, a set of well-distributed solutions over the entire Pareto front is not always the best solution set. For example, it is reported in a recent study (Ishibuchi et al. (2018)) that the best solution set for IGD minimization has much smaller diversity than a fully expanded solution set over the entire Pareto front when a large number of uniformly distributed reference points are used for IGD calculation for many-objective problems.

In this section, we examine the optimal distribution of solutions for hypervolume maximization using SMS-EMOA (Emmerich et al. (2005); Beume et al. (2007)) for DTLZ1-4 (Deb et al. (2002)) and Minus-DTLZ1-4 (Ishibuchi et al. (2017c)). The objective space of each test problem is normalized for uniform visualization so that the ideal and nadir points are $(0, \dots, 0)$ and $(1, \dots, 1)$, respectively. Since DTLZ2-4 have the same Pareto front, they have the same optimal distribution of solutions. For the same reason, Minus-DTLZ2-4 have the same optimal distribution of solutions.

In order to focus on the search for the optimal distribution of solutions on the Pareto front, the number of distance variables in each test problem is specified as zero in our computational experiments in this section. Under this setting, all feasible solutions of each test problem are Pareto optimal. SMS-EMOA with the SBX crossover (crossover probability: 1.0, distribution index: 20) and the polynomial mutation (mutation probability: $1/n$ where n is the number of decision variables, distribution index: 20) is applied to each test problem using the following parameter specifications: The population size is 50, and the termination condition is 100,000 generations. Five settings of the reference point $\mathbf{r} = (r, \dots, r)$ are examined in SMS-EMOA: $r = 1.0$ (the same as the nadir point), $r = 1.1$ (10% larger than the nadir point), $r = 1.5$ (50% larger), $r = 2.0$ (100% larger) and $r = 10$ (ten times larger). During the execution of SMS-EMOA, the reference point is fixed at the pre-specified point (i.e., one of the five settings). For each

setting of the reference point, SMS-EMOA is executed five times. In this manner, we obtained 25 solution sets. Each solution set is evaluated using the hypervolume indicator with each setting of the reference point. Then, the best solution set with respect to the hypervolume indicator is selected among the 25 solution sets as the final result for each setting of the reference point.

2.1 Three-Objective DTLZ1 and Minus-DTLZ1 Test Problems

The obtained solution set for each setting of the reference point is shown in Figure 1 for the three-objective normalized DTLZ1. When the nadir point is used as the reference point in Figure 1 (a), the three extreme solutions of the Pareto front are not included in the best solution set. In Figure 1 (b)-(e) with $r = 1.1, 1.5, 2.0, 10$, the same solution set is selected independent of the reference point specification. The solution set in Figure 1 (a) with $r = 1.0$ is similar to the solution set in Figure 1 (b)-(e). These observations suggest that a set of well-distributed solutions over the entire Pareto front is highly evaluated by the hypervolume indicator, independent of the reference point specification. As a result, it is likely that similar results are obtained from hypervolume-based comparison, independent of the reference point specification for the three-objective DTLZ1.

In Figure 2, we show the obtained solution sets for the three-objective normalized Minus-DTLZ1. When the nadir point is used as the reference point in Figure 2 (a), all solutions are inside the Pareto front. On the contrary, when $r = 10$ (i.e., when the reference point is far away from the nadir point) in Figure 2 (e), almost all solutions are on the sides of the Pareto front. Only a single inside solution is included in the best solution set in Figure 2 (e). By increasing the distance of the reference point from the nadir point, the optimal distribution of solutions is biased toward the sides of the Pareto front. Figure 2 clearly shows that the optimal distribution of solutions strongly depends on the location of the reference point. This means that hypervolume-based comparison results for the three-objective Minus-DTLZ1 strongly depend on the reference point specification. Different specifications of the reference point may lead to different comparison results of EMO algorithms by the hypervolume indicator.

The question is why our simulation results in Figure 1 and Figure 2 are totally




Figure 1: Obtained solution sets for the three-objective normalized DTLZ1.




Figure 2: Obtained solution sets for the three-objective normalized Minus-DTLZ1.

different from each other. First, we explain the reason for the difference between Figure 1 (a) and Figure 2 (a) where the nadir point is used as the reference point (i.e., $r = 1.0$). In this case, the hypervolume calculation is performed within the triangular pyramid stretched from the nadir point $(1, 1, 1)$ as shown in Figure 3 and Figure 4. In Figure 3, the triangular Pareto front of DTLZ1 is inscribed in the green inverted triangle, which is the base of the triangular pyramid for the hypervolume calculation. Only the three extreme solutions of the Pareto front are on the sides of the green inverted triangle (i.e., only these extreme solutions have zero hypervolume contribution). Thus, Pareto optimal solutions on the sides of the Pareto front have positive hypervolume contributions. As a result, some solutions on the sides of the Pareto front are included in the best solution set in Figure 1 (a). However, in Figure 4, the green inverted triangle perfectly overlaps the inverted triangular Pareto front of Minus-DTLZ1. This means that any solution on the sides of the Pareto front cannot have a positive hypervolume contribution. As a result, all solutions in the best solution set in Figure 2 (a) are inside the Pareto front.

Next, let us discuss the difference between Figure 1 (e) and Figure 2 (e) when the reference point is far away from the nadir point. This difference can be explained by projecting two types of Pareto fronts onto a two-dimensional subspace as shown in Figure 5. In Figure 5 (a), the triangular Pareto front of DTLZ1 is projected to the red



Figure 3: Hypervolume calculation from the nadir point for the three-objective normalized DTLZ1.



Figure 4: Hypervolume calculation from the nadir point for the three-objective normalized Minus-DTLZ1.



(a) Triangular Pareto front.

(b) Inverted triangular Pareto front.

Figure 5: Projection of the Pareto front to the f_2 - f_3 space.

triangle in the f_2 - f_3 space. Only the extreme solution $(1, 0, 0)$ projected to $(0, 0)$ is non-dominated in the f_2 - f_3 space. This means that only this solution has a positive hypervolume contribution in the f_2 - f_3 space. By moving the reference point away from the Pareto front along the f_1 axis, the hypervolume contribution of only this solution increases in the original three-dimensional objective space. From the same discussion in the f_1 - f_2 and f_1 - f_3 spaces, we can see that the hypervolume contributions of the three extreme solutions increase by moving the reference point far away from the nadir point. The location of the reference point has no effect on the hypervolume contributions of all the other solutions when the three extreme solutions are included in the solution set. Thus the same best solution set is obtained in Figure 1 (b)-(e).

In Figure 5 (b), the inverted triangular Pareto front of Minus-DTLZ1 is projected to the red triangle in the f_2 - f_3 space. All solutions on the line between $(1, 0)$ and $(0, 1)$ are non-dominated in the f_2 - f_3 space. This means that all solutions on this line have positive hypervolume contributions in the f_2 - f_3 space. By moving the reference point away from the Pareto front along the f_1 axis, their hypervolume contributions monotonically increase in the original three-dimensional objective space. From the same discussion in the f_1 - f_2 and f_1 - f_3 spaces, we can see that the hypervolume contributions of all solutions on the sides of the inverted triangular Pareto front monotonically increase with the increase in the distance of the reference point from the nadir point. As a result, more solutions on the sides of the Pareto front are included in the best solution set in Figure 2. When r is very large, almost all solutions in the best solution set are on the sides of the Pareto front as in Figure 2 (e). In Section 3, we will explain these discussions about the relation between the location of the reference point and the optimal solution set through visualization of the hypervolume contribution of each solution.

2.2 Five-Objective DTLZ1 and Minus-DTLZ1 Test Problems

Experimental results on the five-objective normalized DTLZ1 and Minus-DTLZ1 are shown in Figure 6 and Figure 7, respectively. In Figure 6, the same solution set is obtained for the three settings of the reference point ($r = 1.5, 2.0, 10$) for the five-objective normalized DTLZ1. This is because the location of the reference point has an effect only on the hypervolume contributions of the five extreme solutions of the five-objective normalized DTLZ1. When $r = 1.0$ (i.e., when the nadir point is used as the reference point), the best solution set does not cover the entire Pareto front. That is, the diversity of the best solution set in Figure 6 (a) is much smaller than that in Figure 6 (c)-(e). The best solution set for $r = 1.1$ does not cover the entire Pareto front, either. That is, $r = 1.1$ is too small for the five-objective normalized DTLZ1 whereas it is a good setting for the three-objective normalized DTLZ1. This observation suggests that the appropriate specification of the reference point depends on the number of objectives. This issue will be further discussed in Section 3.

In Figure 7, a different solution set is obtained for each setting of the reference point for the five-objective Minus-DTLZ1. When $r = 1.0$, the diversity of the obtained solution set is very small in Figure 7 (a). Almost all solutions are close to the center of the Pareto front $(0.8, 0.8, 0.8, 0.8, 0.8)$. When $r = 1.1$, the diversity is still small in Figure 7 (b) whereas it is larger than that in Figure 7 (a). However, when $r = 10$, no solutions are close to the center of the Pareto front in Figure 7 (e). Our experimental results in Figure 6 and Figure 7 suggest that the reference point specification is much more important for Minus-DTLZ1 in Figure 7 than DTLZ1 in Figure 6. This is consistent with our observation in Figure 1 and Figure 2 for the three-objective normalized DTLZ1 and Minus-DTLZ1.




Figure 6: Obtained solution sets for the five-objective normalized DTLZ1 problem.




Figure 7: Obtained solution sets for the five-objective Minus-DTLZ1 problem.

2.3 Three-Objective DTLZ2 and Minus-DTLZ2 Test Problems

Whereas DTLZ1 and Minus-DTLZ1 have linear Pareto fronts, the Pareto fronts of DTLZ2 and Minus-DTLZ2 are non-linear. Experimental results on the three-objective DTLZ2 and normalized Minus-DTLZ2 are shown in Figure 8 and Figure 9, respectively. As in Figure 1 and Figure 6 for DTLZ1, the location of the reference point does not have a large effect on the optimal distribution of solutions in Figure 8 for DTLZ2. This is because DTLZ2 has a triangular Pareto front. In Figure 9 for Minus-DTLZ2, the location of the reference point has a large effect on the optimal distribution of solutions. This is because Minus-DTLZ2 has an inverted Triangular Pareto front. However, the optimal




Figure 8: Obtained solution sets for the three-objective DTLZ2 problem.




Figure 9: Obtained solution sets for the three-objective Minus-DTLZ2 problem.




Figure 10: Hypervolume contribution of each of the five uniformly distributed solutions.

distributions of solutions in Figure 9 for Minus-DTLZ2 are clearly different from Figure 2 for Minus-DTLZ1. For example, in Figure 2 (e), almost all solutions are on the sides of the inverted triangular Pareto front. However, most solutions are not on the sides of the inverted triangular Pareto front in Figure 9 (e) whereas they are close to the sides. This is related to the nonlinear shape of the Pareto front of Minus-DTLZ2.

As explained in Auger et al. (2012) for a nonlinear Pareto front of a two-objective problem, the two extreme solutions of the Pareto front are not always included in the optimal distribution of solutions even when the reference point is far away from the Pareto front. In Figure 10, we show the hypervolume contribution of each of the five uniformly distributed solutions on the entire Pareto fronts of the two-objective DTLZ2 and normalized Minus-DTLZ2. The hypervolume contribution of each solution is shown by the corresponding shaded rectangle in Figure 10 (a) and (b), and by the size of the corresponding open circle in Figure 10 (c) and (d). The size of each open circle in Figure 10 (c) and (d) is proportional to the hypervolume contribution of the corresponding solution. This hypervolume visualization method is used for three-objective problems later in this paper. When the reference point is far away from the Pareto front, the two extreme solutions of DTLZ2 have large hypervolume contributions as shown in Figure 10 (a) and (c). Thus they are included in the optimal distribution of solutions of DTLZ2. However, the two extreme solutions of Minus-DTLZ2 have small hypervolume contributions as shown in Figure 10 (b) and (d). Thus they are not included in the optimal distribution of solutions of Minus-DTLZ2. Similar explanations hold for the three-objective normalized Minus-DTLZ2 in Figure 9. Thus most solutions in the obtained best solution set are inside the Pareto front (whereas they are close to the sides of the Pareto front) even when the reference point is far away from the Pareto front.

In this section, we demonstrated that the optimal distribution of solutions for hypervolume maximization depends on the location of the reference point, the curvature property of the Pareto front (linear, convex, or concave) and the Pareto front shape (triangular or inverted triangular). Among the six combinations of the curvature property and the Pareto front shape, four combinations are examined using DTLZ1 (linear triangular), Minus-DTLZ1 (linear inverted triangular), DTLZ2 (concave triangular) and Minus-DTLZ2 (concave inverted triangular).

3 Reference Point Specification Method

In this section, we propose a reference point specification method in hypervolume calculation for fair performance comparison of EMO algorithms. First, we derive a simple specification method for two-objective problems from theoretical analysis of the opti-

mal distribution of solutions on the linear Pareto front of a normalized two-objective problem. Next, the specification method is generalized for multi-objective problems. Then, its validity is examined through computational experiments.

3.1 Basic Idea for Two-Objective Problems: Theoretical Analysis

Let us consider a two-objective minimization problem with a linear Pareto front, which is a line between $(0, 1)$ and $(1, 0)$ in the two-dimensional objective space. For this problem, the hypervolume indicator is maximized by a set of equidistant points on the Pareto front (Emmerich et al. (2007); Auger et al. (2012)). The optimal distribution of five solutions is illustrated in Figure 11 for different settings of the reference point $r = (r, r)$: $r = 1.0, 1.1, 1.25, 1.5$. Each shaded region in Figure 11 shows the hypervolume contribution of the corresponding solution. As illustrated in Figure 11 (a) with $r = 1.0$, the two extreme solutions of the Pareto front are not included in the optimal solution set when the nadir point is used as the reference point. When $r = 1.1$ in Figure 11 (b), they are not included in the optimal solution set either. The two extreme solutions are included in the optimal solution set only when the reference point satisfies the inequality condition $r \geq 1 + 1/(\mu - 1)$ where μ is the number of solutions. This condition has already been proven in Theorem 3 of Brockhoff (2010). In Figure 11 with $\mu = 5$, this condition means $r \geq 1.25$. Thus the two extreme solutions are included in Figure 11 (c) and (d).

In Figure 12, we show the hypervolume contribution of each of the five uniformly distributed solutions. When the nadir point is used as the reference point in Figure 12 (a), the two extreme solutions have no hypervolume contribution. This means that they have no effect on hypervolume-based performance comparison results. This is not desirable for fair performance comparison. When $r = 1.1$ in Figure 12 (b), the




Figure 11: Optimal distributions of solutions for hypervolume maximization for a linear Pareto front.




Figure 12: Hypervolume contribution of each of the uniformly distributed solutions on a linear Pareto front.

two extreme solutions have much smaller hypervolume contributions than the other three solutions. However, when $r = 1.5$ in Figure 12 (d), they have much larger hypervolume contributions than the other three solutions. All of these settings are not desirable from a viewpoint of fair performance comparison since some solutions in the solution set have much smaller or much larger effects on hypervolume calculation (i.e., on hypervolume-based performance comparison results). Only when $r = 1.25$ (i.e., $r = 1 + 1/(\mu - 1)$), all solutions in Figure 12 have the same hypervolume contribution. Based on these discussions, we propose to specify the reference point $r = (r, r)$ as $r = 1 + 1/(\mu - 1)$ for two-objective problems.

The optimal distribution of solutions has been actively studied theoretically for the case of two objectives. For example, an interesting relation between the optimal distribution and the hypervolume contribution was proven by Bader (2009): Each solution in the optimal distribution for hypervolume maximization has the same hypervolume contribution when the number of solutions increases to infinity (Theorem 3.26). It was shown by Auger et al. (2012) that the density of solutions in the optimal distribution depends only on the slope of the Pareto front. Generalization of the hypervolume indicator was also proposed. For example, cone-based hypervolume indicators were proposed in Emmerich et al. (2013) where the optimal distribution of solutions depends on an angle parameter of cones. When the angle parameter is small (i.e., cones are acute), solutions are more evenly distributed on convex and concave Pareto fronts than the case of the standard hypervolume indicator. Reference point free weighted hypervolume indicators were proposed in Emmerich et al. (2014) where it was shown that the standard hypervolume indicator belongs to a more general class of Pareto compliant measures. Whereas those generalized versions were proposed, we focus on the standard hypervolume indicator since it has frequently been used for performance comparison of EMO algorithms in the literature.

3.2 Generalization to Multi-Objective Problems: Proposed Method

In Figure 13, we show 15 uniformly distributed solutions on the entire Pareto front of the three-objective normalized DTLZ1. In this paper, uniformly distributed solutions are generated by the simplex lattice-based method in the same manner as MOEA/D for generating uniformly distributed weight vectors (whereas other methods may generate other distributions, e.g., simplex centroid). In Figure 13, each side of the triangular Pareto front is divided into four intervals as in Figure 11 (c) and Figure 12 (c) with five solutions on the linear Pareto front. Our idea is to apply the reference point specification method in the previous subsection to the sides of the triangular Pareto front. Since each side is divided into four intervals in Figure 13 as in Figure 11 (c) and Figure 12 (c), we propose to use $(1.25, 1.25, 1.25)$ as the reference point in Figure 13. In general, let




Figure 13: Fifteen uniformly distributed solutions for the three-objective normalized DTLZ1 problem.

us assume that each side of the triangular Pareto front of the normalized m -objective DTLZ1 problem is divided into H intervals. Then, we propose to specify the reference point $\mathbf{r} = (r, r, \dots, r)$ as $r = 1 + 1/H$. In the case of two-objective problems, H is calculated from the number of solutions μ as $H = \mu - 1$. Thus the proposed specification method by $r = 1 + 1/H$ can be viewed as a generalization of our basic idea (i.e., $r = 1 + 1/(\mu - 1)$) for two-objective problems in the previous subsection.

In MOEA/D (Zhang and Li (2007)), the population size μ is defined for an m -objective problem using a user-defined integer parameter H as $\mu = C_{m-1}^{H+m-1}$ where C_m^n is the number of m -combinations from a set of n elements, i.e., $C_m^n = n!/m!(n-m)!$. The integer parameter H in MOEA/D corresponds to the number of intervals of each side of the triangular Pareto front (e.g., $H = 4$ in Figure 13). When the population size μ is specified by this formulation, we can directly use our formulation (i.e., $r = 1 + 1/H$) for specifying the reference point $\mathbf{r} = (r, r, \dots, r)$. However, in general, the population size μ is not always specified as $\mu = C_{m-1}^{H+m-1}$. For handling such a general case, first we specify the value of H using the following formulation:

$$C_{m-1}^{H+m-1} \leq \mu < C_{m-1}^{H+m}, \quad (1)$$

where μ is the population size and m is the number of objectives. Then, the reference point $\mathbf{r} = (r, r, \dots, r)$ is specified in the normalized objective space with the ideal point $(0, 0, \dots, 0)$ and the nadir point $(1, 1, \dots, 1)$ as follows:

$$r = 1 + 1/H. \quad (2)$$

Our proposal is to use (1) and (2) for specifying the reference point $\mathbf{r} = (r, r, \dots, r)$ in the normalized objective space from the population size μ and the number of objectives m .

3.3 Numerical Examples: Hypervolume Contribution of Each Solution on Linear Pareto Fronts

In this subsection, we examine whether each of the uniformly distributed solutions over the entire Pareto front has a similar hypervolume contribution when the reference point $\mathbf{r} = (r, r, \dots, r)$ is specified by the proposed DTL method. In Figure 14, we generate 66 solutions for the three-objective normalized DTLZ1 using the weight vector generation method in MOEA/D with $H = 10$. The suggested reference point by the proposed method is $r = 1.1$. The hypervolume contribution of each solution is shown by the size of the corresponding circle. That is, the area of each circle is proportional to the hypervolume contribution of the corresponding solution. In this subsection, the size of each circle is normalized so that the largest circle in each figure has the same maximum radius. So, we can compare the size of each circle within each figure (e.g., between a circle in Figure 14 (a) and another one in Figure 14 (e)). However, we cannot compare the size of each circle between different figures (e.g., between Figure 14 and Figure 15).

As we have already explained, the hypervolume contributions of the three extreme solutions are zero when the nadir point is used as the reference point (i.e., $r = 1.0$) in Figure 14 (a). When r is large (e.g., $r = 2.0$), those three extreme solutions have much larger hypervolume contributions than the other solutions. Only when the reference point is specified by the proposed method as $r = 1.1$, all the 66 solutions have the same hypervolume contribution.

For examining the effect of the number of solutions on the appropriate reference point specification, we also generate 21 solutions using the weight vector generation




Figure 14: Hypervolume contribution of each solution for the three-objective normalized DTLZ1 ($H = 10$).




Figure 15: Hypervolume contribution of each solution for the three-objective normalized DTLZ1 ($H = 5$).

method in MOEA/D with $H = 5$. The suggested reference point by the proposed method is $r = 1.2$. The hypervolume contribution of each solution is shown in Figure 15 in the same manner as in Figure 14 (i.e., the area of each circle is proportional to the hypervolume contribution of the corresponding solution). As we have already explained, the size of each circle is normalized in each figure. So, the three extreme solutions in Figure 14 (e) and Figure 15 (e) have the largest circles of the same size. However, their hypervolume contributions in Figure 15 are larger than those in Figure 14.

When $r = 1.1$ in Figure 15 (b), the three extreme solutions have much smaller hypervolume contributions than the other solutions. This observation suggests that the reference point is too close to the Pareto front. When $r = 1.5$ and $r = 2.0$, the three extreme solutions have much larger hypervolume contributions than the other solutions. This observation suggests that those specifications are too far from the Pareto front. Only when the reference point is specified by the proposed method as $r = 1.2$, all the 21 solutions have the same hypervolume contribution.

We also perform the same computational experiment for the three-objective normalized Minus-DTLZ1 for the case of $H = 5$ (i.e., 21 solutions). Experimental results are shown in Figure 16. Only in this figure, the three extreme solutions which are shown by squares (instead of circles) in Figure 16 (e) are excluded from the normalization of the size of each circle. This is because they have too large hypervolume contributions to be visualized in the same figure.

Minus-DTLZ1 has an inverted triangular Pareto front. When the nadir point is used as the reference point in Figure 16 (a), all solutions on the sides of the Pareto front have no hypervolume contribution. When $r = 1.1$, those solutions have much smaller hypervolume contribution than the other solutions inside the Pareto front. Only when we specify the reference point as $r = 1.2$ using the proposed method, all the 21 solutions have the same hypervolume contribution. By increasing the value of r (i.e., by moving




Figure 16: Hypervolume contribution of each solution for the three-objective normalized Minus-DTLZ1 ($H = 5$).




Figure 17: Hypervolume contribution of each solution for the five-objective normalized DTLZ1 ($H = 2$). The hypervolume contribution is shown by the line thickness.

the reference point away from the Pareto front), the hypervolume contributions of the three extreme points increase as in the case of the triangular Pareto front in Figure 15. However, at the same time, the hypervolume contributions of the other solutions on the sides of the inverted triangular Pareto front also increase in Figure 16. This is the reason why almost all solutions in the best solution sets are on the sides of the inverted triangular Pareto front when the reference point is far away from the Pareto front (e.g., see Figure 9 (e)). It should be noted that the hypervolume contributions of those solutions do not change by moving the location of the reference point in the case of the triangular Pareto front as shown in Figure 15. That is, when the Pareto front is triangular, the location of the reference point has an effect only on the hypervolume contributions of the extreme solutions.

To examine the effect of the number of objectives on the appropriate reference point specification, we generate 15 solutions for the five-objective normalized DTLZ1 using the weight vector generation method in MOEA/D with $H = 2$. The suggested reference point by the proposed method is $r = 1.5$. The hypervolume contribution of each solution is shown by the line thickness in Figure 17. When $r = 1.0$, the five extreme solutions with $f_i = 1$ for one of the five objectives have no hypervolume contribution. Thus they are not depicted in Figure 17 (a). When $r = 1.1$ and $r = 1.2$, their hypervolume contributions are much smaller than the other solutions. When the suggested reference point $r = 1.5$ is used in Figure 17 (d), all solutions have the same hypervolume contribution. By increasing the value of r from $r = 1.5$, the five extreme solutions have larger hypervolume contributions than the other solutions as shown in Figure 17 (e).

We further examine the hypervolume contributions of a large number of solutions of the five-objective normalized DTLZ1. First, we generate its 1001 solutions uniformly on the triangular Pareto front using the weight vector generation mechanism of MOEA/D with $H = 10$. The suggested reference point by the proposed method is 1.1.

Table 1: Hypervolume contribution of each solution of the five-objective normalized DTLZ1 ($\times 10^{-5}$)

Solution Type (Number of solutions)	$r = 1.0$	$r = 1.05$	$r = 1.1$	$r = 1.2$	$r = 1.5$
One non-zero element (5)	0.000	0.500	1.000	2.000	5.000
Two non-zero elements (90)	1.000	1.000	1.000	1.000	1.000
Three non-zero elements (360)	1.000	1.000	1.000	1.000	1.000
Four non-zero elements (420)	1.000	1.000	1.000	1.000	1.000
Five non-zero elements (126)	1.000	1.000	1.000	1.000	1.000

Table 2: Hypervolume contribution of each solution of the five-objective normalized Minus-DTLZ1 ($\times 10^{-5}$)

Solution Type (Number of solutions)	$r = 1.0$	$r = 1.05$	$r = 1.1$	$r = 1.2$	$r = 1.5$
One non-one element (5)	0.000	0.0625	1.000	16.00	625.0
Two non-one elements (90)	0.000	0.125	1.000	8.000	125.0
Three non-one elements (360)	0.000	0.250	1.000	4.000	25.00
Four non-one elements (420)	0.000	0.500	1.000	2.000	5.000
Five non-one elements (126)	1.000	1.000	1.000	1.000	1.000

The 1001 solutions (i.e., 1001 points in the five-dimensional objective space) satisfy the following relations: $f_1 + f_2 + f_3 + f_4 + f_5 = 1$ and $0 \leq f_i \leq 1$ for $i = 1, 2, \dots, 5$. Among them, 126 solutions are inside the Pareto front. That is, they have no zero element (i.e., they have five non-zero elements). All the others are boundary solutions with at least one zero element (i.e., with less than five non-zero elements). For example, the five extreme solutions have only a single non-zero element: $(1, 0, 0, 0, 0)$, $(0, 1, 0, 0, 0)$, $(0, 0, 1, 0, 0)$, $(0, 0, 0, 1, 0)$ and $(0, 0, 0, 0, 1)$. Ninety solutions have only two non-zero elements such as $(0.9, 0.1, 0, 0, 0)$ and $(0, 0, 0.4, 0, 0.6)$. The hypervolume contribution of each solution is summarized in Table 1. When the reference point is specified by the proposed method as $r = 1.1$, all solutions have exactly the same hypervolume contribution. The hypervolume contributions of only the five extreme solutions depend on the location of the reference point in Table 1. This is because the shape of the Pareto front is triangular as we have already explained.

In the same manner as in Table 1, we also examine the hypervolume contribution of 1001 uniformly distributed solutions on the inverted triangular Pareto front of the five-objective normalized Minus-DTLZ1. The suggested reference point by the proposed method is 1.1. The 1001 solutions satisfy the following relations: $f_1 + f_2 + f_3 + f_4 + f_5 = 4$ and $0 \leq f_i \leq 1$ for $i = 1, 2, \dots, 5$. Among the 1001 solutions, 126 are inside the Pareto front. Those inside solutions have five non-one elements. All the others are boundary solutions with less than five non-one elements such as the five extreme solutions with only a single non-one element: $(0, 1, 1, 1, 1)$, $(1, 0, 1, 1, 1)$, $(1, 1, 0, 1, 1)$, $(1, 1, 1, 0, 1)$ and $(1, 1, 1, 1, 0)$. The hypervolume contribution of each solution is summarized in Table 2. When the reference point is specified by the proposed method as $r = 1.1$, all solutions have exactly the same hypervolume contribution. When the nadir point is used as the reference point (i.e., $r = 1.0$), only the 126 inside solutions have positive hypervolume contributions. Their hypervolume contributions are the same independent of the loca-




Figure 18: Hypervolume contribution of each solution for the three-objective DTLZ2 ($H = 5$).




Figure 19: Hypervolume contribution of each solution for the three-objective normalized Minus-DTLZ2 ($H = 5$).

tion of the reference point in Table 2. The hypervolume contribution of each of the other boundary solutions increases as the value of r increases. It is interesting to observe that the increase of the hypervolume contribution is more rapid when solutions have less non-one elements. For example, the hypervolume contributions of the five extreme solutions with only a single non-one element drastically increase when the value of r increases from $r = 1.1$ to $r = 1.5$ in Table 2.

3.4 Numerical Examples: Hypervolume Contribution of Each Solution on Nonlinear Pareto Fronts

Experimental results in the previous subsection show that each solution in a uniformly distributed solution set on a linear Pareto front has the same hypervolume contribution when the reference point $\mathbf{r} = (r, r, \dots, r)$ is specified by the proposed method as $r = 1 + 1/H$. For comparison, we also examine the hypervolume contribution of each solution on non-linear Pareto fronts. In Figure 18, we generate 21 solutions for the three-objective DTLZ2 using the weight vector generation method in MOEA/D with $H = 5$. More specifically, we calculate the best solution for each weight vector with respect to the PBI function with $\theta = \infty$ in MOEA/D. Then, we calculate the hypervolume contribution of each solution. When the nadir point is used as the reference point (i.e., $r = 1.0$) in Figure 18 (a), the three extreme solutions have no hypervolume contribution. When $r = 2.0$ in Figure 18 (e), their hypervolume contributions are much larger than the other solutions. Due to the nonlinearity of the Pareto front, each solution has a different hypervolume contribution in Figure 18. The proposed specification $r = 1.2$ in Figure 18 (c) is more appropriate than $r = 1.0$ and $r = 2.0$ in the sense that the difference of the hypervolume contributions among the 66 solutions is small. Due to the concave shape of the Pareto front, solutions on the sides of the Pareto front have larger hypervolume contributions than inside solutions in Figure 18 as explained for the two-objective DTLZ2 in Figure 10.

We also generate 21 solutions for the three-objective normalized Minus-DTLZ2

using the inverted PBI function with $\theta = \infty$ in MOEA/D (Sato (2014)). Then, we calculate the hypervolume contribution of each solution. When the nadir point is used as the reference point (i.e., $r = 1.0$) in Figure 19 (a), all solutions on the sides of the Pareto fronts have no hypervolume contribution. By increasing the distance of the reference point from the nadir point, their hypervolume contributions increase. However, due to the convex shape of the Pareto front, solutions on the sides of the Pareto front have smaller hypervolume contributions than inside solutions in Figure 19 as explained for the two-objective Minus-DTLZ2 in Figure 10.

3.5 Numerical Examples: Hypervolume-based Comparison of Solution Sets

In this subsection, we examine whether a uniformly distributed solution set has a larger hypervolume than biased solution sets when the reference point is specified by the proposed method. In our computational experiments, first we generate a set of 91 uniformly distributed solutions for the three-objective normalized DTLZ1 by specifying the value of H as $H = 12$ in the weight vector specification mechanism in MOEA/D as shown in Figure 20 (d). For comparison, we also generate other solution sets with 91 solutions. For generating a biased solution set around the center of the Pareto front, first we generate 136 uniformly distributed solutions for $H = 15$. This solution set consists of 91 inside solutions and the other 45 solutions on the sides of the Pareto front. We use a set of the 91 inside solutions for comparison as shown in Figure 20 (c). We also generate a further biased solution set. First, we generate 190 solutions by specifying H as $H = 18$. Next, we remove the 54 solutions on the sides of the Pareto front. Then, we remove the 45 solutions on the side of the remaining solution set. We use the other 91 inside solutions as a solution set for comparison as shown in Figure 20 (b). In the same manner, we also generate a solution set with 91 solutions selected from 253 uniformly distributed solutions for $H = 21$ in Figure 20 (a).

We also generate biased solution sets toward the sides of the Pareto front in the following manner. First, we generate a set of 66 uniformly distributed solutions by specifying the value of H as $H = 10$. This solution set includes 36 inside solutions. In addition to those 36 inside solutions, 55 solutions are uniformly distributed over the sides of the Pareto front to generate a solution set with 91 solutions as shown in Figure 20 (e). In the same manner, we generate a solution set from $H = 8$ in Figure 20 (f) where the 21 inside solutions are generated from $H = 8$ and the other 70 solutions are uniformly distributed over the sides of the Pareto front. We also generate a solution set with 10 inside solutions from $H = 6$ and 81 uniformly distributed solutions on the sides of the Pareto front in Figure 20 (g).

In the same manner, we also generate seven solution sets with 91 solutions of the three-objective normalized Minus-DTLZ1, DTLZ2, and Minus-DTLZ2. The generated




Figure 20: Generated solution sets with 91 solutions of the three-objective normalized DTLZ1 for hypervolume-based performance comparison. The value of H shows the resolution of inside solutions. A large value of H means a dense distribution of solutions around the center of the Pareto front.




Figure 21: Generated solution sets with 91 solutions of the three-objective normalized Minus-DTLZ2.

Table 3: Best solution set by the hypervolume-based comparison with each specification of the reference point.

Test Problem	Reference Point Specification (r)										
	1.01	1.02	1.04	1.06	1.08	1.1	1.2	1.4	1.6	1.8	2
DTLZ1	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)
Minus-DTLZ1	(c)	(c)	(c)	(d)	(d)	(d)	(e)	(e)	(f)	(f)	(f)
DTLZ2	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)	(d)
Minus-DTLZ2	(b)	(b)	(b)	(b)	(b)	(b)	(c)	(c)	(d)	(d)	(d)

solution sets of Minus-DTLZ2 are shown in Figure 21. The generated seven solution sets for each test problem are compared by the hypervolume indicator for each of the following specifications of the reference point: $r = 1.00, 1.02, 1.04, 1.06, 1.08, 1.10, 1.20, 1.40, 1.60, 1.80, 2.00$. Table 3 shows the solution set with the largest hypervolume for each specification of the reference point. For DTLZ1, the well-distributed solution set in Figure 20 (d) is evaluated as the best solution set, independent of the specification of the reference point. For DTLZ2, such a solution set is also evaluated as the best solution set, independent of the specification of the reference point. However, for Minus-DTLZ1 and Minus-DTLZ2, the hypervolume-based performance comparison results strongly depend on the specification of the reference point. The suggested reference point by the proposed method for a three-objective problem with 91 solutions (i.e., $m = 3$ and $\mu = 91$) is $r = 1 + 1/12 = 1.0833$. In Table 3, the well-distributed solution set over the entire Pareto front of Minus-DTLZ1, which is the inverted distribution of Figure 20 (d), is evaluated as the best solution set only when $r = 1.06, 1.08, 1.10$. This observation supports the validity of the proposed reference point specification method. However, for Minus-DTLZ2, the solution set in Figure 21 (b) is evaluated as the best solution set when $r = 1.06, 1.08, 1.10$. This observation, which is consistent with the above-mentioned experimental results on Minus-DTLZ2 with respect to the best solution set and the hypervolume contribution of each solution, suggests the necessity of special care in the hypervolume-based performance comparison for multi-objective problems with convex Pareto fronts.

4 Computational Experiments on Various Test Problems

We have already explained that an appropriate reference point can be obtained by the proposed method for DTLZ1, DTLZ2 and Minus-DTLZ1. We have also demonstrated that the location of the reference point has a large effect on the optimal distribution of solutions and performance comparison results through computational experiments on Minus-DTLZ1 and Minus-DTLZ2 with inverted triangular Pareto fronts. However, the effect of the location of the reference point is small when test problems have triangular

Pareto fronts (e.g., DTLZ1-4 and WFG4-9). In this section, we examine the effect of the reference point specification using other frequently-used three-objective test problems: knapsack problem, car-side impact problem, and WFG3. The number of solutions is specified as 50. For $m = 3$ (three objectives) and $\mu = 50$ (50 solutions), the suggested reference point by the proposed method is calculated as $r = 1.125$ since H is determined as $H = 8$ from Eq. (1). The three test problems in this section are normalized as three-objective minimization problems with the ideal point $(0, 0, 0)$ and the nadir point $(1, 1, 1)$.

4.1 Three-Objective Knapsack Problem

Multi-objective knapsack problems have frequently been used as combinatorial multi-objective test problems in the EMO community (Zitzler and Thiele (1999); Zhang and Li (2007); Ishibuchi et al. (2015)). As a test problem, we use the three-objective 500-item knapsack problem in Ishibuchi et al. (2015). First, we search for an approximate Pareto front by MOEA/D with Tchebycheff function (Zhang and Li (2007)) using a large computation load (population size: 10,011, termination condition: 1,000 generations, execution: ten times). All solutions obtained from ten runs of MOEA/D are merged into a single solution set. Next, we select only non-dominated solutions from the merged solution set. In this manner, we obtain an approximate Pareto front with 31,509 non-dominated solutions in Figure 22 (a) where the three-dimensional objective space is normalized as a minimization problem with the ideal point $(0, 0, 0)$ and the nadir point $(1, 1, 1)$. Then, a greedy algorithm (Guerreiro et al. (2016)) is used to select a set of 50 solutions for hypervolume maximization. In the greedy algorithm, the first solution is selected by examining the hypervolume indicator of each solution. That is, the best solution with the largest hypervolume is selected as the first solution and included in the solution set. Next, the second solution is included in the solution set to maximize the hypervolume indicator after examining all combinations of the first solution and one of the other solutions. Then, the third solution is included in the solution set to maximize the hypervolume indicator after examining all combinations of the first two solutions and one of the other solutions. In this manner, 50 solutions are selected for each of the following settings of the reference point: $r = 1.01, 1.125, 1.5, 2.0$. The obtained solution sets are shown in Figure 22 (b)-(e).

When the reference point is slightly worse than the nadir point (i.e., $r = 1.01$) in Figure 22 (a), only a few solutions are close to the sides of the Pareto front. When the reference point is far away from the Pareto front (i.e., $r = 2.0$) in Figure 22 (e), many solutions are close to the sides of the Pareto front. When the reference point is specified by the proposed method (i.e., $r = 1.125$) in Figure 22 (c), solutions are




Figure 22: Pareto front in (a) and selected 50 solutions for each specification of the reference point for hypervolume maximization in (b)-(e) for the three-objective 500-item knapsack problem.

well distributed over the entire Pareto front. Our experimental results in Figure 22 show that an appropriate reference point is specified by the proposed method for the three-objective 500-item knapsack problem. They also show that the optimal solution set strongly depends on the reference point specification. That is, hypervolume-based performance comparison results are sensitive to the reference point specification. As shown in Figure 22, the Pareto front of the three-objective knapsack problem is inverted triangular. As a result, the experimental results in Figure 22 are similar to those on the three-objective Minus-DTLZ1 problem in Figure 2.

4.2 Three-objective Car-Side Impact Problem

The car-side impact problem has frequently been used as a multi-objective problem with constraint conditions in the EMO community (Jain and Deb (2014); Asafuddoula et al. (2015)). This problem has three objectives, ten constraint conditions and eleven decision variables. First, we search for an approximate Pareto front by NSGA-II (Deb et al. (2002)) using a large computation load (population size: 5,000, termination condition: 10,000 generations, execution: 20 times). In NSGA-II, a set of 5,000 feasible solutions are generated as an initial population by randomly generating a large number of solutions and selecting only feasible solutions. During the execution of NSGA-II, infeasible solutions are handled as being dominated by feasible ones. All solutions obtained from 20 runs of NSGA-II are merged into a single solution set, from which non-dominated solutions are selected. In this manner, we obtain an approximate Pareto front with 54,874 non-dominated solutions in Figure 23 (a) in the three-dimensional objective space. Then, a greedy algorithm is used to select a set of 50 solutions for hypervolume maximization.

In Subsection 4.1, we used MOEA/D to search for an approximate Pareto front of the three-objective knapsack problem. This is because much better results were reported for multi-objective knapsack problems with 2-4 objectives from MOEA/D than NSGA-II in the literature (e.g., Zhang and Li (2007); Ishibuchi et al. (2015)). However, in this subsection, we used NSGA-II to search for an approximate Pareto front of the car-side impact problem. This is because each objective of the car-side impact problem has a different scale. Since MOEA/D has no scaling mechanism of the objective space, usually it does not work well on such a multi-objective problem (e.g., WFG test problems of Huband et al. (2006)).

The selected 50 solutions are shown in Figure 23 (b)-(e). When a reference point is slightly worse than the nadir point (i.e., $r = 1.01$) in Figure 23 (b), no solutions are close to the right-hand side boundary of the Pareto front. By specifying the reference point by the proposed method as $r = 1.125$, well-distributed solutions are obtained in Figure 23 (c) except for the right-bottom corner of the Pareto front. By increasing the value of




Figure 23: Pareto front in (a) and selected 50 solutions for each specification of the reference point for hypervolume maximization in (b)-(e) for the car-side impact problem.




Figure 24: Pareto front in (a) and obtained 50 solutions for each specification of the reference point for hypervolume maximization in (b)-(e) for the three-objective WFG3.

r (i.e., by moving the reference point far away from the Pareto front), more solutions are obtained around the top and the right sides of the Pareto front as shown in Figure 23 (d) and (e). Our experimental results on the car-side impact problem show that an appropriate reference point is specified by the proposed method. They also show that the optimal solution set strongly depends on the reference point specification. That is, hypervolume-based performance comparison results are sensitive to the reference point specification. The shape of the Pareto front of the car-side impact problem is a mixture of triangle and inverted triangle. In Figure 23, the bottom-left half of the Pareto front is triangular. The solution at the bottom-right corner is the single optimal solution in the $f_2 - f_3$ space as in Figure 5 (a). So, many solutions are not around the bottom and left sides of the Pareto front even when $r = 2.0$ in Figure 23 (e). On the contrary, the top-right half of the Pareto front is inverted triangular. So, many solutions are around the top and right sides of the Pareto front in Figure 23 (e).

4.3 Three-Objective WFG3 Problem

WFG3 was originally proposed as a degenerate test problem (Huband et al. (2006)). However, it was shown in Ishibuchi et al. (2016) that WFG3 has a partially degenerate Pareto front as shown in Figure 24 (a). In the same manner as in Section 2, SMS-EMOA is applied to the three-objective WFG3. The number of distance variables is specified as two due to the restriction of the WFG problem formulation whereas it was specified as zero for DTLZ1 and DTLZ2 in Section 2. Experimental results are shown in Figure 24 (b)-(e). When $r = 1.01$ in Figure 24 (b), the two extreme solutions at the bottom-left and bottom-right corners are not selected. When $r = 2.0$ in Figure 24 (e), only four solutions are inside the Pareto front. If compared with these results, the solution set from $r = 1.125$ in Figure 24 (c) looks better. As in Figure 22 and Figure 23, Figure 24 shows that an appropriate reference point is specified by the proposed method. Figure 24 also shows that the best solution set depends on the reference point specification. That is, hypervolume-based performance comparison results depend on the reference point specification.

5 Effect on the Search Ability of SMS-EMOA

In this section, we examine whether the proposed specification method is useful in SMS-EMOA or not. In Sections 2-4, we discussed the reference point specification method for fair performance comparison where the true or approximate Pareto front is assumed to be known. However, in the use of the proposed method in SMS-EMOA, the ideal and nadir points are estimated from the current population. Their estimations are not accurate especially in early generations. In our computational experiments in this section, the normalization is performed using non-dominated solutions among so-

lutions in the current population at each generation. As mentioned in Emmerich et al. (2005), SMS-EMOA can work with an infinite reference point. This is a good specification for two-objective problems and multi-objective problems with triangular Pareto fronts. However, this is not appropriate for multi-objective problems with inverted triangular Pareto fronts. In this section, we examine the usefulness of the proposed reference point specification method in SMS-EMOA for multi-objective problems with triangular and inverted triangular Pareto fronts.

We apply SMS-EMOA to the three-objective normalized DTLZ1 and Minus-DTLZ1 under the following parameter specifications:

- Number of distance variables: 5,
- Population size (μ): 15 and 50,
- Crossover: SBX with the index 20 (probability: 1.0),
- Mutation: Polynomial mutation with the index 20 (probability: $1/n$ where n is the number of variables),
- Termination conditions: 100,000 generations,
- Number of runs: 5.

In our computational experiments, we examine the following three-specification of the reference point: a slightly worse reference point than the nadir point (i.e., $r = 1.01$), the suggested reference point (i.e., $r = 1.25$ for $\mu = 15$ and $r = 1.125$ for $\mu = 50$), and a much worse reference point than the nadir point (i.e., $r = 2.0$).

Experimental results on DTLZ1 are shown in Figures 25-27 where the final solution sets obtained from the five runs are shown for each specification of the population size.




Figure 25: Obtained solution sets for the three-objective normalized DTLZ1 with $r = 1.01$ (top: population size 15, bottom: population size 50).




Figure 26: Obtained solution sets for the three-objective normalized DTLZ1 with the suggested reference point (top: population size 15 and $r = 1.25$, bottom: population size 50 and $r = 1.125$).




Figure 27: Obtained solution sets for the three-objective normalized DTLZ1 with $r = 2.0$ (top: population size 15, bottom: population size 50).

When $r = 1.01$ in Figure 25, good results are not obtained. The final population always converges to a small region of the Pareto front. When the reference point is specified by the proposed method in Figure 26, well-distributed solutions are always obtained. Good results are also obtained in Figure 27 where the reference point is far away from the Pareto front. As we have already explained, the reference point can be far away from the triangular Pareto front. However, such a reference point specification has a clear negative effect when the Pareto front is inverted triangular.

Experimental results on Minus-DTLZ1 are shown in Figures 28-30. When the reference point is far away from the Pareto front in Figure 30, many solutions are on the sides of the inverted triangular Pareto front. When the reference point is slightly worse than the nadir point (i.e., $r = 1.01$), the final population always converges to a small region




Figure 28: Obtained solution sets for the three-objective normalized Minus-DTLZ1 with $r = 1.01$ (top: population size 15, bottom: population size 50).




Figure 29: Obtained solution sets for the three-objective normalized Minus-DTLZ1 with the suggested reference point (top: population size 15 and $r = 1.25$, bottom: population size 50 and $r = 1.125$).




Figure 30: Obtained solution sets for the three-objective normalized Minus-DTLZ1 with $r = 2.0$ (top: population size 15, bottom: population size 50).

of the Pareto front in Figure 28. When the reference point is specified by the proposed method in Figure 29, well-distributed solutions are obtained. Our experimental results in Figures 28-30 show that the performance of SMS-EMOA on multi-objective problems with inverted triangular Pareto fronts strongly depends on the reference point specification. Only when the reference point is appropriately specified, well-distributed solutions over the entire Pareto front are obtained.

We also apply SMS-EMOA to the five-objective normalized DTLZ1 and Minus-DTLZ1. The population size is specified as 50. The suggested reference point for this setting (i.e., $m = 5$ and $\mu = 50$) is $r = 1.333$. Experimental results are shown in Figure 31 and Figure 32. When the reference point is slightly worse than the nadir point (i.e., $r = 1.01$), good results are not obtained for these test problems as shown in the top five results in each figure. When the reference point is specified by the proposed method (i.e., $r = 1.333$), good results are obtained as shown in the middle five results in each figure. When the reference point is far away from the Pareto front (i.e., $r = 2.0$), good solution sets are obtained for DTLZ1 in the bottom five results of Figure 31. This is because DTLZ1 has a triangular Pareto front. However, in the application of SMS-EMOA to Minus-DTLZ1, most solutions are around the sides of the Pareto front (i.e., at least one objective value is 0 or 1). Many solutions are not obtained inside the Pareto front (whereas this is not very clear in the bottom five results in Figure 32).




Figure 31: Experimental results on the five-objective normalized DTLZ1 with the population size 50 (top: $r = 1.01$, middle: $r = 1.333$, bottom: $r = 2.0$).




Figure 32: Experimental results on the five-objective normalized Minus-DTLZ1 with the population size 50 (top: $r = 1.01$, middle: $r = 1.333$, bottom: $r = 2.0$).

6 Concluding Remarks

In this paper, we discussed the reference point specification in hypervolume calculation for fair performance comparison of EMO algorithms. First, we demonstrated that the optimal distribution of solutions for hypervolume maximization strongly depends on the location of the reference point except for multi-objective problems with triangular Pareto fronts such as DTLZ1-4 and WFG4-9. This means that hypervolume-based performance comparison results strongly depend on the location of the reference point. It was shown through computational experiments that a slightly worse reference point than the nadir point is not always appropriate especially for the case of many-objective problems and/or small population size. It was also shown that a much worse reference point, which is far away from the nadir point, is not appropriate except for multi-objective problems with triangular Pareto fronts. Based on the theoretical analysis of the optimal distribution of solutions for hypervolume maximization, we proposed a reference point specification method. The proposed method calculates the location of the reference point based on the number of solutions (i.e., population size) and the number of objectives. In the proposed method, the distance between the reference point and the nadir point increases with the increase in the number of objectives and the decrease in the number of solutions (i.e., it increases with the increase in the distance between adjacent solutions). Our experimental results showed that an appropriate reference point can be specified by the proposed method for multi-objective and many-objective problems with linear triangular and linear inverted triangular Pareto fronts. It was also shown that the reference point specified by the proposed method is more appropriate for other test problems than a slightly or much worse reference point.

Since the proposed method was derived from the theoretical analysis of the optimal distribution of solutions for two-objective problems with linear Pareto fronts, the specified reference point is not necessarily appropriate for multi-objective problems with nonlinear Pareto fronts. Modification of the proposed method for nonlinear Pareto fronts is an interesting future research topic. Since the proposed method was derived for fair performance comparison where the true Pareto front is assumed to be known, the specified reference point is not necessarily appropriate in hypervolume-based EMO algorithms where the ideal and nadir points are estimated from the current population. Modification of the proposed method for hypervolume-based EMO algorithms is also an interesting future research topic. Especially, on-line adaptation of the reference point

seems to be a promising future research direction.

In this paper, we discussed the reference point specification in hypervolume calculation using the optimal distribution of solutions for hypervolume maximization. This approach (i.e., use of the optimal distribution of solutions) applies to the analysis of other performance indicators (Ishibuchi et al. (2018)). Comparison of performance indicators using the optimal distribution of solutions for each indicator is an important future research direction for their further understanding and fair performance comparison of EMO algorithms.

In this paper, we implicitly assumed that the size of each solution set to be compared was the same. That is, the number of non-dominated solutions obtained by each EMO algorithm was the same. However, this is not always the case. As demonstrated in Ishibuchi et al. (2016), an appropriate population size for each EMO algorithm is different. As a result, performance comparison results of EMO algorithms depend on the setting of the population size when they are compared under the same population size. Moreover, even when they are compared under the same population size, the number of obtained non-dominated solutions by each EMO algorithm may be different since all the obtained solutions are not always non-dominated. Thus, fair performance comparison of EMO algorithms is difficult. In order to address this difficulty, two scenarios were proposed in Ishibuchi et al. (2016) and used in Tanabe et al. (2017). One scenario is the use of an unbounded archive where all non-dominated solutions among examined ones are stored. Performance comparison is based on all non-dominated solutions in the obtained archive by each run of an EMO algorithm. The other scenario is the selection of a pre-specified number of non-dominated solutions from the obtained archive. By using several settings of the number of non-dominated solutions, performance comparison results may be unbiased if compared with the case of the common population size.

Acknowledgment

This work was supported by the Science and Technology Innovation Committee Foundation of Shenzhen (Grant No. ZDSYS201703031748284).

References

- Asafuddoula, M., Ray, T., and Sarker, R. (2015). A decomposition-based evolutionary algorithm for many objective optimization. *IEEE Transactions on Evolutionary Computation*, 19(3):445–460.
- Auger, A., Bader, J., Brockhoff, D., and Zitzler, E. (2012). Hypervolume-based multiobjective optimization: Theoretical foundations and practical implications. *Theoretical Computer Science*, 425:75–103.
- Bader, J. (2009). *Hypervolume-Based Search for Multiobjective Optimization: Theory and Methods*. PhD thesis, ETH Zurich.
- Bader, J. and Zitzler, E. (2011). HypE: An algorithm for fast hypervolume-based many-objective optimization. *Evolutionary Computation*, 19(1):45–76.
- Basseur, M., Derbel, B., Goffon, A., and Liefoghe, A. (2016). Experiments on greedy and local search heuristics for d -dimensional hypervolume subset selection. In *2016 Genetic and Evolutionary Computation Conference (GECCO 2016)*, pages 541–548.
- Beume, N., Naujoks, B., and Emmerich, M. (2007). SMS-EMOA: Multiobjective selection based on dominated hypervolume. *European Journal of Operational Research*, 181(3):1653–1669.

- Bringmann, K., Friedrich, T., and Klitzke, P. (2014). Two-dimensional subset selection for hypervolume and epsilon-indicator. In *2014 Genetic and Evolutionary Computation Conference (GECCO 2014)*, pages 589–596.
- Brockhoff, D. (2010). Optimal μ -distributions for the hypervolume indicator for problems with linear bi-objective fronts: Exact and exhaustive results. In *8th International Conference on Simulated Evolution and Learning (SEAL 2010)*, pages 24–34.
- Chugh, T., Jin, Y., Miettinen, K., Hakanen, J., and Sindhya, K. (2018). A surrogate-assisted reference vector guided evolutionary algorithm for computationally expensive many-objective optimization. *IEEE Transactions on Evolutionary Computation (Early Access Paper: Available from IEEE Xplore)*.
- Deb, K., Thiele, L., Laumanns, M., and Zitzler, E. (2002). Scalable multi-objective optimization test problems. In *2002 IEEE Congress on Evolutionary Computation (CEC 2002)*, pages 825–830.
- Emmerich, M., Beume, N., and Naujoks, B. (2005). An EMO algorithm using the hypervolume measure as selection criterion. In *2005 International Conference on Evolutionary Multi-Criterion Optimization (EMO 2005)*, pages 62–76.
- Emmerich, M., Deutz, A., and Beume, N. (2007). Gradient-based/evolutionary relay hybrid for computing Pareto front approximations maximizing the S-metric. In *4th International Workshop on Hybrid Metaheuristics (HM 2007)*, LNCS 4771, pages 140–156.
- Emmerich, M., Deutz, A., Krusselbrink, J., and Shukla, P. K. (2013). Cone-based hypervolume indicators: Construction, properties, and efficient computation. In *2013 International Conference on Evolutionary Multi-Criterion Optimization (EMO 2013)*, pages 111–127.
- Emmerich, M., Deutz, A., and Yevseyeva, I. (2014). On reference point free weighted hypervolume indicators based on desirability functions and their probabilistic interpretation. *Procedia Technology*, 16:532–541.
- Friedrich, T., Neumann, F., and Thyssen, C. (2015). Multiplicative approximations, optimal hypervolume distributions, and the choice of the reference point. *Evolutionary Computation*, 23(1):131–159.
- Guerreiro, P., Fonseca, C. M., and Paquete, L. (2015). Greedy hypervolume subset selection in the three-objective case. In *2015 Genetic and Evolutionary Computation Conference (GECCO 2015)*, pages 671–678.
- Guerreiro, P., Fonseca, C. M., and Paquete, L. (2016). Greedy hypervolume subset selection in low dimensions. *Evolutionary Computation*, 24(3):521–544.
- Huband, S., Hingston, P., Barone, L., and While, L. (2006). A review of multiobjective test problems and a scalable test problem toolkit. *IEEE Transactions on Evolutionary Computation*, 10(5):477–506.
- Ishibuchi, H., Akedo, N., and Nojima, Y. (2015). Behavior of multi-objective evolutionary algorithms on many-objective knapsack problems. *IEEE Transactions on Evolutionary Computation*, 19(2):264–283.
- Ishibuchi, H., Imada, R., Setoguchi, Y., and Nojima, Y. (2017a). Hypervolume subset selection for triangular and inverted triangular Pareto fronts of three-objective problems. In *14th ACM/SIGEVO Conference on Foundations of Genetic Algorithms (FOGA 2017)*, pages 95–110.
- Ishibuchi, H., Imada, R., Setoguchi, Y., and Nojima, Y. (2017b). Reference point specification in hypervolume calculation for fair comparison and efficient search. In *2017 Genetic and Evolutionary Computation Conference (GECCO 2017)*, pages 585–592.
- Ishibuchi, H., Imada, R., Setoguchi, Y., and Nojima, Y. (2018). Reference point specification in inverted generational distance for triangular linear Pareto front. *IEEE Transactions on Evolutionary Computation (Early Access Paper: Available from IEEE Xplore)*.

- Ishibuchi, H., Masuda, H., and Nojima, Y. (2014). Selecting a small number of non-dominated solutions to be presented to the decision maker. In *2014 IEEE International Conference on Systems, Man, and Cybernetics (SMC 2014)*, pages 3850–3855.
- Ishibuchi, H., Setoguchi, Y., Masuda, H., and Nojima, Y. (2016). How to compare many-objective algorithms under different settings of population and archive sizes. In *2016 IEEE Congress on Evolutionary Computation (CEC 2016)*, pages 1149–1156.
- Ishibuchi, H., Setoguchi, Y., Masuda, H., and Nojima, Y. (2017c). Performance of decomposition-based many-objective algorithms strongly depends on Pareto front shapes. *IEEE Transactions on Evolutionary Computation*, 21(2):169–190.
- Jain, H. and Deb, K. (2014). An evolutionary many-objective optimization algorithm using reference-point based non-dominated sorting approach, Part II: Handling constraints and extending to an adaptive approach. *IEEE Transactions on Evolutionary Computation*, 18(4):602–622.
- Knowles, J. D., Corne, D. W., and Fleischer, M. (2003). Bounded archiving using the Lebesgue measure. In *2003 IEEE Congress on Evolutionary Computation (CEC 2003)*, pages 2490–2497.
- Kuhn, T., Fonseca, C. M., Paquete, L., Ruzika, S., Duarte, M. M., and Figueira, J. R. (2016). Hypervolume subset selection in two dimensions: Formulations and algorithms. *Evolutionary Computation*, 24(3):411–425.
- Li, K., Deb, K., Zhang, Q., and Kwong, S. (2015). An evolutionary many-objective optimization algorithm based on dominance and decomposition. *IEEE Transactions on Evolutionary Computation*, 19(5):694–716.
- Maltese, J., Ombuki-Berman, B. M., and Engelbrecht, A. P. (2018). A scalability study of many-objective optimization algorithms. *IEEE Transactions on Evolutionary Computation (Early Access Paper: Available from IEEE Xplore)*.
- Sato, H. (2014). Inverted PBI in MOEA/D and its impact on the search performance on multi and many-objective optimization. In *2014 Genetic and Evolutionary Computation Conference (GECCO 2014)*, pages 645–652.
- Seada, H. and Deb, K. (2016). A unified evolutionary optimization procedure for single, multiple, and many objectives. *IEEE Transactions on Evolutionary Computation*, 20(3):358–369.
- Shukla, P. K., Doll, N., and Schmeck, H. (2014). A theoretical analysis of volume based Pareto front approximations. In *2014 Genetic and Evolutionary Computation Conference (GECCO 2014)*, pages 1415–1422.
- Tanabe, R., Ishibuchi, H., and Oyama, A. (2017). Benchmarking multi- and many-objective evolutionary algorithms under two optimization scenarios. *IEEE Access*, 5:19597–19619.
- Wagner, T., Beume, N., and Naujoks, B. (2007). Pareto-, aggregation-, and indicator-based methods in many-objective optimization. In *4th International Conference on Evolutionary Multi-Criterion Optimization (EMO 2007)*, pages 742–756.
- Yuan, Y., Xu, H., Wang, B., and Yao, X. (2016a). A new dominance relation based evolutionary algorithm for many-objective optimization. *IEEE Transactions on Evolutionary Computation*, 20(1):16–37.
- Yuan, Y., Xu, H., Wang, B., Zhang, B., and Yao, X. (2016b). Balancing convergence and diversity in decomposition-based many-objective optimizers. *IEEE Transactions on Evolutionary Computation*, 20(2):180–198.
- Zhang, Q. and Li, H. (2007). MOEA/D: A multiobjective evolutionary algorithm based on decomposition. *IEEE Transactions on Evolutionary Computation*, 11(6):712–731.
- Zhang, X., Tian, Y., Cheng, R., and Jin, Y. (2018). A decision variable clustering based evolutionary algorithm for large-scale many-objective optimization. *IEEE Transactions on Evolutionary Computation (Early Access Paper: Available from IEEE Xplore)*.

- Zitzler, E., Brockhoff, D., and Thiele, L. (2007). The hypervolume indicator revisited: On the design of Pareto-compliant indicators via weighted integration. In *4th International Conference on Evolutionary Multi-Criterion Optimization (EMO 2007)*, pages 862–876.
- Zitzler, E. and Thiele, L. (1998). Multiobjective optimization using evolutionary algorithms - a comparative case study. In *5th International Conference on Parallel Problem Solving from Nature (PPSN V)*, pages 292–301.
- Zitzler, E. and Thiele, L. (1999). Multiobjective evolutionary algorithms: A comparative case study and the strength Pareto approach. *IEEE Transactions on Evolutionary Computation*, 3(4):257–271.
- Zitzler, E., Thiele, L., Laumanns, M., Fonseca, C. M., and da Fonseca, V. G. (2003). Performance assessment of multiobjective optimizers: An analysis and review. *IEEE Transactions on Evolutionary Computation*, 7(2):117–132.